

# Flat Point 2016

## Fine Scale Rocky Shore Monitoring



Prepared for  
Greater  
Wellington  
Regional  
Council  
April  
2016

Cover Photo: Flat Point - looking south toward Glenburn headland.



Flat Point - looking north towards the sand-spit and subtidal reef system.

# **Flat Point 2016**

## **Fine Scale Rocky Shore Monitoring**

**Prepared for  
Greater Wellington Regional Council**

**By**

**Ben Robertson and Leigh Stevens**

### **RECOMMENDED CITATION**

*Robertson, B.P. and Stevens, L.M. 2016. Flat Point 2016 Fine Scale Rocky Shore Monitoring. Prepared by Wriggle Coastal Management for Greater Wellington Regional Council. 16p. plus appendices.*

# Contents

Rocky Shore - Executive Summary . . . . .	vii
1. Introduction . . . . .	1
2. Methods . . . . .	5
3. Results and Discussion . . . . .	7
4. Summary . . . . .	15
5. Monitoring . . . . .	16
6. Acknowledgements . . . . .	16
7. References . . . . .	16
Appendix 1. Detailed Results . . . . .	17

## List of Figures

Figure 1. Location of the rocky shore sampling area at Flat Point . . . . .	4
Figure 2. General shoreline zonation present at Flat Point . . . . .	5
Figure 3. <i>Chamaeosipho columna</i> and <i>Xenostrobus neozelanicus</i> on the high shore . . . . .	9
Figure 4. <i>Sypharochiton pelliserpentis</i> and <i>Cellana radians</i> on the mid shore . . . . .	10
Figure 5. <i>Hormosira banksii</i> and <i>Corallina officinalis</i> on the low shore . . . . .	10
Figure 6. Topshells, limpets and chitons from the upper, mid and low eu littoral zones . . . . .	11
Figure 7. n-MDS plot showing the relationship in community composition for Flat Point, Jan. 2016. . . . .	12
Figure 8. Mean SACFOR rating for species present in 6 fixed quadrats in high, mid and low eu littoral zones. . . . .	14

## List of Tables

Table 1. Summary of the major environmental issues affecting NZ rocky shores. . . . .	2
Table 2. SACFOR Percentage Cover and Density Scales (after Marine Nature Conservation Review - MNCR) . . . . .	6
Table 3. Results of the semi-quantitative SACFOR assessment at Flat Point, 24-25 January 2016 . . . . .	8
Table 4. Summary of richness, abundance and diversity indices in quadrats, Flat Point, 2016. . . . .	12
Table 5. Raw data, mean number/percentage cover, and SACFOR rating of shore quadrats, Flat Point, 2016. . . . .	13

All photos by Wriggle except where noted otherwise.



# ROCKY SHORE - EXECUTIVE SUMMARY



This report summarises results of the first year of fine scale monitoring of the rocky shore community at Flat Point, Wairarapa, an eastern coast site exposed to moderate-high wave energy, northerly, easterly and southerly winds, and bathed by the relatively cool waters of the Wairarapa Coastal Current. It is a key site in Greater Wellington Regional Council's (GWRC's) long-term coastal monitoring programme. This report describes:

- Fine scale semi-quantitative monitoring of the abundance and diversity of conspicuous rocky shore plants and animals, and
- Fine scale quantitative monitoring of the abundance and diversity of plants and animals in 18 x 0.25m<sup>2</sup> fixed quadrats at high, mid, and low eulittoral (intertidal) levels at three sites.

## FINE SCALE MONITORING RESULTS

The semi-quantitative monitoring identified 27 attached or sessile rocky shore species, excluding those in heavily fissured areas and rock pools. In total, 11 algae, 3 limpet, 7 topshell, 1 chiton, 3 barnacle and 2 bivalve species were observed.

The uppermost shore was dominated by eroding coastal duneland fringing a sandy beach, with supralittoral (splash zone) rocky reef habitat limited to the tops of large rocks/bedrock exposed above the beach sands. Present on the rocks were topshells (3), including the brown and blue-banded periwinkles (superabundant/rare), 2 limpets (common/rare), 3 barnacles (superabundant/rare), the black mussel (rare), and the red algae *Porphyra* sp. (frequent).

The high eulittoral (intertidal) zone [10 species] was characterised by a dense cover of barnacles - column barnacles (superabundant), 2 limpet species (abundant/frequent), 3 topshell species (frequent/rare), snakeskin chiton (common), *Porphyra* sp. (occasional), and blue and black mussels (frequent/rare).

The mid shore [10 species] was largely dominated by column barnacles (abundant). Also present were 2 limpet species (common/occasional), 4 topshells (common/rare), snakeskin chiton (common), and 2 bivalves (rare).

The lower shore [17 species] was most diverse and was dominated by a superabundant cover of turfing, crustose red algae, and an abundant cover of brown algae (Nephture's necklace). The algal canopy provided shelter and refuge to a range of other species, including: limpets (3), barnacles (1), chitons (1), topshells (2), and subdominant brown (5), red (2) and green (1) algae.

Nineteen of the 27 species recorded in the semi-quantitative monitoring were also recorded in the first baseline year of quantitative fixed quadrat sampling.

## ROCKY SHORE ISSUES AND CONDITION

Rocky shore ecology on the Wairarapa coast has been assessed as having a low-moderate risk, primarily due to predicted climate change pressures including accelerated sea level rise, temperature change, ocean acidification and, to a lesser extent, over-collection of living resources and introduction of invasive species. The risk from pathogens, sedimentation, eutrophication, and toxins was considered low.

The first year of baseline monitoring found the coastline in a healthy and unpolluted condition. No introduced invasive species were seen, and there was no indication of excessive nutrient or sediment inputs.

## RECOMMENDED MONITORING AND MANAGEMENT


To provide a robust baseline of rocky shore conditions (particularly to enable monitoring of changes from predicted accelerated sea level rise and increased temperatures), it is recommended that a three year annual fine scale monitoring baseline be established, and then monitored 5 yearly thereafter. The next monitoring is scheduled for January 2017.

While the rocky shore baseline is established, it is proposed that condition ratings be developed to characterise the status of the shore. It is proposed that these ratings focus on measuring shifts in community composition, the presence or absence of key indicator species (including introduced plants and animals), and indicators of nutrient enrichment and sedimentation.



# 1. INTRODUCTION

## OVERVIEW



Developing an understanding of the condition and risks to coastal habitats is critical to the management of biological resources. Importantly, the “Wairarapa Coastal Habitats - Mapping, Risk Assessment and Monitoring” report (Robertson and Stevens 2007) identified a low-moderate risk to rocky shore ecology on the Wairarapa coast. This was primarily from predicted climate change effects of accelerated sea level rise and elevated temperatures, the over-collection of living resources, and the introduction of invasive species. The primary ecological responses to such pressures are considered to be habitat change, and effects on biodiversity. Due to the generally high clarity, low nutrients, and low disease risk of water that bathes the Wairarapa rocky shoreline, the risk from pathogens, sedimentation, eutrophication, and toxins was considered low. Because of this, the number of monitoring indicators can be kept small, although this may change if catchment land use intensifies.

Therefore, to address the identified risks, and to provide baseline information on rocky shore ecology at key representative locations, Robertson and Stevens (2007) recommended long term monitoring of the abundance and diversity of plants and animals at regionally representative high diversity rocky shores (e.g. Flat Point and Cape Palliser) using rapid assessment methods developed under the Marine Biodiversity and Climate Change Project (Hiscock 1996). Wriggle Coastal Management was contracted by Greater Wellington Regional Council (GWRC) to undertake the first year of a 3 year baseline of annual monitoring at Flat Point in 2016. After establishment of the baseline, monitoring is scheduled to be undertaken 5 yearly with the results used to help to determine the extent to which the coast is affected by major environmental pressures (Table 1), both in the short and long term.

Rocky shores are a dominant and visually dramatic part of the Wairarapa coastline. They reflect the erosive effect of waves where softer rocks are worn down, leaving harder rocks exposed. The habitat is physically complex, with rockpools, gullies, crevices and boulders providing a diverse range of habitats supporting a variety of different species. The harsh and variable physical conditions, including light availability, degree of exposure, large shifts in temperature and salinity, aspect, substrate, and biotic features, lead to the development of a characteristic zonation of species on stable shoreline substrate. These include zones dominated by lichens, periwinkles, barnacles, limpets, mussels, and canopy forming algae - the dominant biogenic habitat along temperate rocky shores worldwide (e.g. Tomanek and Helmuth 2002).

Canopy forming algae play a vital role on the rocky shore by providing food and shelter to a wide range of species. Consequently, any change or loss of this canopy habitat is likely to result in a cascade of related effects. For example, canopy loss will increase heat stress, desiccation of understory species, and wave exposure, likely resulting in a simplified cover dominated by resilient species e.g. coralline algae, which in turn may preclude the re-establishment of canopy species. Changes in canopy cover may also result in secondary impacts altering existing ecosystem dynamics, with bare space colonised by new species (possibly invasive or nuisance species), food shortages altering grazing dynamics or predation, or changed susceptibility to other stressors such as sedimentation and eutrophication.

The relationship between stressors (both natural and human influenced) and changes to rocky shore communities is complex and can be highly variable. However, there are clear links between the degradation of rocky shore habitat and the combined effects of elevated nutrient, sediment, pathogen, and toxin inputs, harvesting, trampling, coastal development, introduced species, as well as broader stressors such as changes to sea temperature, sea level, wave exposure, and storm frequency and intensity (directly influenced by global climate change) - see Table 1.

As such, monitoring representative rocky shore sites provides a robust and effective way of detecting changes to this important and highly valued coastal community.

# 1. Introduction (Continued)

**Table 1. Summary of the major environmental issues affecting NZ rocky shores.**

There are five main environmental issues that affect NZ rocky shores, with the main stressors being climate change and sea level rise, over-collection of living resources, introduction of invasive species, and pollution. All these can be linked to a decline in the dominant algal canopy species, on which many other species depend for food or habitat:

## 1. Habitat Loss or Modification.

**Climate Change and Sea level Rise.** Predicted climate change impacts (e.g. warmer temperatures, ocean acidification, sea-level rise, increased storm frequency) are expected to alter species ranges (e.g. increased sub-tropical introductions and/or establishment of pest species), alter planktonic and kelp production, and interfere with the formation of shells and skeletons by corals, crabs, marine snails, and bivalves. Long term predictions are the loss of rare species, a reduction in species diversity, and the loss of entire communities of organisms in some situations.

**Over-collection of Living Resources and Recreation.** Direct removal of living resources (e.g. fish, mussels, paua, crayfish, algae) can cause major community level changes (e.g. Airoldi et al. 2005) from disruption to natural predator-prey balances or loss of habitat-maintaining species. For example, some popular recreational fish species (e.g. greenbone, red moki) play an important role in maintaining algal habitat and depletion of these species can cause significant changes in community structure (e.g. Taylor and Schiel 2010). Macroalgal harvesting can remove protective habitat, resulting in species loss and greater exposure to natural disturbances. Impacts are expected from recreational activities (e.g. algal trampling) and over-collection at both local and regional scales, and is likely to intensify as expanding human populations put further pressure on resources.

**Introduction of Invasive Species.** Increased global transport (hull fouling and ballast water discharges) is a major vector in the introduction of invasive or pest plants and animals. Displacement of native species, particularly following disturbance events (e.g. canopy loss), can result in less diverse communities and possibly increased ephemeral blooms. Introduced toxic microalgae, while harmless enough at low levels, can reproduce explosively when conditions are right, giving rise to toxic algal blooms (TABs), and resultant illness and/or mortality of humans, fish, sea birds and marine mammals that ingest toxic fish or shellfish poisoned by TABs. Significant effort and cost may be needed to remove or prevent the spread of unwanted species e.g. *Undaria* - an introduced golden brown seaweed that has been a prominent marine pest in New Zealand with extensive effort put into both minimising its spread removing it from certain locations e.g. Fiordland, Stewart Island.

## 2. Disease Risk.

If pathogen inputs to the coastal area are excessive (e.g. from coastal wastewater discharges or proximity to a contaminated river plume), the disease risk from bathing, wading or eating shellfish can increase to unacceptable levels. High flushing and dilution mean disease risk is unlikely to be significant away from point source discharges. Public health reports of illness are likely to be the first indication of faecal bacterial issues directly impacting on human values and uses.

## 3. Sediment.

Excessive suspended sediments can lower water clarity and cause ecological damage at the shoreline through reduced plant and algal production, clogging of respiratory and suspension feeding organs of sensitive organisms, and can variously affect the ability of recruits to settle and establish (e.g. Airoldi 2003, Foster and Schiel 2010). Sheltered rocky shore habitats, e.g. rockpools, are more susceptible to direct deposition and reduced sediment oxygenation. Generally high wave energy on the open coast will favour offshore sediment settlement over intertidal deposition. Increased sedimentation is likely to reduce biodiversity through lowered productivity and recruitment success, and reduced ability to recover from disturbances. Human values and uses will be reduced directly by poor clarity (swimming/diving), and indirectly through biodiversity changes.

## 4. Eutrophication.

Eutrophication occurs when nutrient inputs are excessive, and can have chronic broad scale impacts over whole coastlines. High nutrients support increased localised nuisance macroalgal growth, and with this, opportunistic grazers. Where dominant, they decrease diversity by excluding or out-competing other species, and can be particularly influential in the colonisation of bare space following disturbance events. Elevated nutrients have also been implicated in a trend of increasing frequency of harmful algal blooms (HABs) which can cause illness in humans and close down shellfish gathering and aquaculture operations. High flushing and dilution on relatively remote exposed rocky shores mean the most likely indicators of eutrophication effects will be increases in nuisance macroalgal growths (e.g. *Ulva*) and phytoplankton blooms, and a subsequent reduction in diversity.

## 5. Toxic Contamination.

If potentially toxic contaminant inputs (e.g. heavy metals, pesticides) are excessive, shoreline biodiversity is threatened and shellfish may be unsuitable for eating. Except for large-scale infrequent discharges such as oil spills, pollution tends mainly to influence embayed coastlines or areas immediately adjacent to outfalls. Increased toxins are unlikely to be a significant issue on NZ's exposed outer coasts but if present, will reduce biodiversity and human values and uses.

## 1. Introduction (Continued)

The Flat Point fine scale rocky shore intertidal monitoring site is located approximately 500m south of the region's main swimming and boat launching beach, and directly south of the airstrip (Figures 1a and 1b). The area is representative of the rocky shoreline on this exposed region of the eastern coast, and is characterised by the following:

- Predominantly exposed, rugged reefs, composed of generally soft, easily erodible sandstones and mudstones.
- Exposure to moderate-high wave energy and southerly, northerly and easterly winds.
- Bathed by the relatively cool, and often nutrient depleted, waters of the Wairarapa Coastal Current that flows northeast along the Wairarapa coast. However, inshore waters are influenced by elevated nutrient, sediment and pathogen loadings from Wairarapa river plumes.
- Dominated beneath low water by the branching, brown macroalgae *Carpophyllum flexuosum* and *Cystophora torlosa* (zig-zag weed), and directly above low water by the beaded, brown macroalga *Hormosira banksii* (neptune's necklace), with barnacles and limpets common above the algal zone.

The sampling area was located on a rough, irregular intertidal reef (~100m x 50m) which was seaward of a relatively wide strip of sandy beach, and part of a wider sequence of connected intertidal and subtidal reefs present on this section of coast (see photo below). Several submerged reefs are present offshore. Although the area is semi-exposed, and is periodically subject to high wind and wave action from the southeast, the sampling site itself is relatively well protected from direct wave action, with the sloping and undulating shoreline further helping to dissipate wave energy. Consequently, it supports a relatively diverse rocky shore community and exhibits strong shoreline zonation (Figure 2). The shoreline zones are relatively narrow (each ~1 vertical metre) reflecting the tidal range at the site (~1.2-2.1m).

The site is not directly or significantly influenced by river plumes, terrestrial discharges (e.g. stormwater, sewage), or structures (e.g. seawalls, wharfs, marine farms). Human use is moderate-high. It is a popular tourist destination, a highly valued recreational paua fishery, and is valued for diving, fishing, and its scenic beauty and bird life. The monitoring sites are considered unlikely to be appreciably affected by recreational fishers or visitors because quadrat locations are discretely marked (unlikely to be noticed), are positioned where direct impacts are unlikely, and do not support species commonly targeted for recreational harvesting e.g. paua and mussels.

The current report describes the methods and results of the first year of the proposed 3 year baseline of rocky shore monitoring at Flat Point.



View south over the Flat Point rocky shore sampling site.

# 1. Introduction (Continued)




Figure 1. Location of the rocky shore sampling area at Flat Point.

## 2. METHODS

The fine scale rocky shore monitoring programme involves measuring the abundance and diversity of conspicuous plants and animals. Monitoring targets the supralittoral zone (the area regularly splashed, but not submerged, by seawater) and the eulittoral (intertidal) zone that extends from the rarely inundated spring high water tide line, to the almost always inundated neap low tide line. Results will be used to evaluate any vertical shift in the zonation pattern associated with climate change, or impacts from introduced species, over-collection of shellfish (e.g. paua, mussels), excessive sediment and nutrient inputs, as well as to provide a baseline for infrequent risks such as oil spills.

Sampling was undertaken by two scientists during relatively calm sea conditions on 24-25 January 2016 when estuary monitoring was being undertaken in the region.

The methodology is based on that used in the UK MarClim - Marine Biodiversity and Climate Change Project (MNCR 1990, Hiscock 1996, 1998), and consists of two parts,

1. A semi-quantitative assessment to develop a checklist of the species present and record their relative abundance across a representative sampling area.
2. Recording the abundance and diversity of plants and animals in 0.25 m<sup>2</sup> fixed quadrats positioned in the spatially largest strata at the site, and stratified within 3 eulittoral tide levels (High, Mid, and Low).

The semi-quantitative assessment was applied by walking over and photographing the wider sampling area, and identifying and recording the relative abundance of all the conspicuous species present from the supralittoral zone to mean low water. At each shore zone a time limit of 60 minutes was used to guide the sampling effort, with extensively shaded areas, rock pools or heavily fissured areas excluded from the assessment. Details were recorded on pre-prepared data sheets that included the range of species likely to be found at the site. In addition, a photographic field guide was used to assist with field identifications.

The abundance of each species was rated using SACFOR categories described in Table 2. The SACFOR assessment preferentially uses percentage cover of two growth types of attached organisms - Crust/ Meadow (e.g. lichen, barnacles, coralline paint), or Massive/Turf (e.g. bull kelp, coralline turf) - Table 2, A. All other individual organisms >5mm in size were counted, with the largest individual organism size used to determine the relevant SACFOR size class rating for each species as detailed in Table 2, B.

The semi-quantitative assessment guided the selection of 18 stratified fixed intertidal quadrats, because true random sampling approaches are not appropriate on a broken rocky shore. The use of fixed quadrats reduces the need for extensive sample replication and minimises spatial variation, while seasonal variation is minimised by scheduling monitoring for the same period each year (January to March). Within the wider sampling area, 3 sites were identified on bedrock which were sheltered from the direct effect of prevailing wind and waves, and to facilitate safe sampling (Figure 2). At each site, 6 quadrats were located, 2 each at high, mid and low tide levels.



Figure 2. General shoreline zonation present at Flat Point (note supratidal zone limited to very top of bedrock reef).

## 2. Methods (Continued)



Quadrats at each shore height had similar physical characteristics (slope, aspect, wave exposure), and were positioned in areas with attached plants or animals as the change to these features is the primary focus of the monitoring. The upper shore true left hand corner of each quadrat was marked for repeat sampling by drilling and fixing a stainless steel bolt in the rock, the site location photographed, and GPS position recorded.

After selecting and marking each quadrat, the following information was recorded:

### High Eulittoral Quadrats

(6 quadrats located ~1m below the top of the barnacle zone)

- Percent cover of all barnacles, mussels, and algae.
- Number of each periwinkle species present (counted from a representative 2cm x 2cm section within each quadrat).
- Number of each limpet or chiton (individuals >10mm) in each 0.25m<sup>2</sup> quadrat.

### Mid Eulittoral Quadrats

(6 quadrats in the middle of the barnacle zone)

- Percent cover of all barnacles, mussels, and algae.
- Number of each limpet or chiton (individuals >10mm) in each 0.25m<sup>2</sup> quadrat.
- Number of each species of snail >5mm in each 0.25m<sup>2</sup> quadrat.

### Low Eulittoral Quadrats

(6 quadrats ~1m above the bottom of the barnacle zone)

- Percent cover of all barnacles, mussels, and algae.
- Number of each limpet or chiton (individuals >10mm) in each 0.25m<sup>2</sup> quadrat.
- Number of each species of snail >5mm in each 0.25m<sup>2</sup> quadrat.

**Table 2. SACFOR Percentage Cover and Density Scales (after Marine Nature Conservation Review - MNCR).**


A. PERCENTAGE COVER		
i. Crust/Meadow	% cover	ii. Massive/Turf
S	>80	-
A	40-79	S
C	20-39	A
F	10-19	C
O	5-9	F
R	1-4	O
-	<1	R

SACFOR Category
S = Super Abundant
A = Abundant
C = Common
F = Frequent
O = Occasional
R = Rare

- Whenever percentage cover can be estimated for an attached species, it should be used in preference to the density scale.
- The massive/turf percentage cover scale should be used for all species except those classified under crust/meadow.
- Where two or more layers exist, for instance foliose algae overgrowing crustose algae, total percentage cover can be over 100%.

### B. DENSITY SCALES

SACFOR size class				Density				
i	ii	iii	iv	0.25m <sup>2</sup> (50x50cm)	1.0m <sup>2</sup> (100x100cm)	10m <sup>2</sup> (3.16x3.16m)	100m <sup>2</sup> (10x10m)	1,000m <sup>2</sup> (31.6x31.6m)
<1cm	1-3cm	3-15cm	>15cm					
S	-	-	-	>2500	>10,000			
A	S	-	-	250-2500	1000-9999	>10,000		
C	A	S	-	25-249	100-999	1000-9999	>10,000	
F	C	A	S	3-24	10-99	100-999	1000-9999	>10,000
O	F	C	A	1-2	1-9	10-99	100-999	1000-9999
R	O	F	C	0-<1	0-<1	1-9	10-99	100-999
-	R	O	F			0-<1	1-9	10-99
-	-	R	O				0-<1	1-9
-	-	-	R					0-<1



### 3. RESULTS AND DISCUSSION

Results of the 24-25 January 2016 fine scale rocky shore monitoring at Flat Point are summarised below in two sections: the semi-quantitative assessment, followed by the fixed quadrat sampling.

The semi-quantitative assessment identified 27 species (Table 3), excluding creviced areas and rock pools. Algae were dominant (11 species), but a wide range of common rocky shore organisms able to withstand the physical rigours of the exposed wave environment were present in a predictable zonation across the shoreline. These included barnacles, limpets, chitons, topshells and bivalve shellfish. Notably, no introduced invasive species were identified.

The zonation is represented within the four key zones assessed, the spray zone of the upper shore (supralittoral), and high, mid, and low intertidal (eulittoral) zones. Within these broad zones, most species comprised two broad categories, those either directly attached to the rock (e.g. barnacles, seaweeds), or sessile species such as limpets and chitons, that are physically adapted to high energy wave conditions (they have a broad base and the ability to cling strongly to the rock), or utilise cracks and depressions in the rock for shelter.

Because this regular zonation of attached and sessile organisms is primarily governed by tidal inundation, monitoring changes to the shore composition provides a very effective way of tracking long-term climate change effects such as predicted accelerated sea level rise or increased temperatures/acidity.

#### Supralittoral Zone

At Flat Point, the extent of the supralittoral zone above the high tide line was restricted to the upper portion of the rocky reef habitat and did not extend further landward because the uppermost shore comprised a sandy beach and eroding dune system (see photo below). Because of this, there was only a small area of supratidal rocky substrata present, and this appears to be relatively frequently inundated by seawater. Consequently, it precluded the presence of lichens, a type of plant that typically inhabits rock and boulders at the upper edge of the spray zone.



However, at the lower edge of the supralittoral fringe, where rock substrata was present, were the brown and blue-banded periwinkles *Austrolittorina cincta* and *Austrolittorina antipodum* (below), classified as common and super-abundant, respectively. These small topshells, while extremely tolerant of the sun, tend to congregate in cracks and fissures in the rock that provide some protection from the elements during the day. Other invertebrate species observed were the column and ridged-surf barnacles *Chamaesipho columna* (super-abundant) and *Elminius placatus* (rare), the ornate and tortoise-shell limpets *Cellana ornata* (common) and *Cellana radians* (rare), and the black bivalve mussel *Xenostrobus zelanicus* (rare). Also present was the red macroalgae *Porphyra sp.*, with an overall abundance rating of frequent.



**Table 3. Results of the semi-quantitative SACFOR assessment at Flat Point, 24-25 January 2016.**

Group and Family		Species	Common name	Scale	Class	Supra	High	Mid	Low
Topshells	Littorinidae	<i>Austrolittorina antipodum</i>	Blue banded periwinkle	#	i	S			
	Littorinidae	<i>Austrolittorina cincta</i>	Brown periwinkle	#	i	C			
	Littorinidae	<i>Risellopsis varia</i>	Periwinkle	#	ii			R	
	Turbinidae	<i>Turbo smaragdus</i>	Cats eye	#	ii			R	C
	Muricidae	<i>Haustrum scobina</i>	Oyster borer	#	ii	R	R	R	
	Trochidae	<i>Diloma aethiops</i>	Grooved topshell	#	ii		F	C	C
	Muricidae	<i>Haustrum haustorium</i>	Black rock shell	#	ii		O		
Limpets	Nacellidae	<i>Cellana ornata</i>	Ornate limpet	#	ii	C	A	C	F
	Nacellidae	<i>Cellana radians</i>	Tortoiseshell limpet	#	ii	R	F	O	O
	Siphonariidae	<i>Siphonaria zelandica</i>		#	ii				O
Chitons	Chitonidae	<i>Sypharochiton pelliserpentis</i>	Snakeskin chiton	#	ii		C	C	C
Barnacles	Catophragmidae	<i>Chamaesipho columna</i>	Column barnacles	%	i	S	S	A	R
	Balanidae	<i>Elminius plicatus</i>	Ridged surf barnacle	%	i	R			
	Balanidae	<i>Epopella plicata</i>	Plicate barnacle	%	i	R			
Red Algae	Bangiaceae	<i>Porphyra</i> sp.	Karengo, Nori	%	ii	F	O		
	Corallinaceae	<i>Corallina officinalis</i>	Pink turf	%	ii				S
	Gelidiaceae	<i>Gelidium caulacanthum</i>		%	ii				C
	Rhodomelaceae	<i>Stictosiphonia arbuscula</i>	Moss weed	%	ii				R
Brown Algae	Adenocystaceae	<i>Adenocystis utricularis</i>	Sea bladders	%	ii				R
	Hormosiraceae	<i>Hormosira banksii</i>	Neptune's necklace	%	ii				A
	Splachnidiaceae	<i>Splachnidium rugosum</i>	Gummy weed	%	ii				F
	Scytothamnaceae	<i>Scytothamnus australis</i>		%	ii				C
	Ralfsiaceae	<i>Ralfsia verrucosa</i>	Tar spot/blood crust	%	ii				R
	Scytosiphonaceae	<i>Colpomenia sinuosa</i>	Oyster thief	%	ii				O
Bivalves	Mytilidae	<i>Mytilus galloprovincialis</i>	Blue mussel	%	i		R	R	
	Mytilidae	<i>Xenostrobus neozelanicus</i>	Black mussel	%	i	R	F	R	
Green Algae	Ulvaceae	<i>Ulva lactuca</i>	Sea lettuce	%	i				O



### 3. Results and Discussion (Continued)

#### High Eulittoral Zone

The high eulittoral zone is submerged for short periods on each tide, and although relatively sheltered, is also exposed to seawater via waves and spray depending on swell and wind angle. Because of its exposed nature, the high shore zone was characterised by extensive patches of bare rock that supported a relatively sparse animal and plant community. Dominating this zone was the sessile, filter-feeding column barnacle *Chamaeosipho columna*, which filter-feeds from the water column at high tide. *C. columna* was superabundant (>80% cover), present in extensive sheets across the rock (see photo below). Other dominant invertebrate species included the highly mobile, herbivorous limpets *Cellana ornata* (abundant) and *Cellana radians* (frequent), and the snakeskin chiton *Sypharochiton pelliserpentis* (listed as common). The blue and black bivalve mussels *Mytilus galloprovincialis* and *Xenostrobus neozelanicus*, which formed patchy colonies amongst the barnacles, were rated as rare/frequent. Also present was the red algae, *Porphyra* sp. - occasional.



Figure 3. *Chamaeosipho columna* and *Xenostrobus neozelanicus* on the high shore.



### 3. Results and Discussion (Continued)



Figure 4. *S. pelleriserpentis* and *C. radians* on the mid shore.

#### Mid Eulittoral Zone

Although still containing bare patches, the mid eulittoral zone was dominated by the barnacle *Chamaeosipho columna*, classified as abundant.

Limpets were the most common of the mobile invertebrates with *Cellana radians* and *C. ornata*, both common/occasional. Other invertebrates included the chiton *Sypharochiton pelleriserpentis* and the grooved topshell *Diloma aethiops*, both common, while the other topshells present (*R. varia*, *T. smaragdus* and *H. scobina*) were rated rare - most present sheltering in shaded cracks and crevices on the vertical rock faces (Figure 4). Also present among the barnacles, were the two species of mussel *M. galloprovincialis* and *X. neozelanicus*, both listed as rare.



Mid Eulittoral

#### Low Eulittoral Zone

The lower eulittoral zone is exposed to the air for only a short period on each tidal cycle and is where algae of various forms have their stronghold on the shore. It was dominated by an extensive (superabundant) close turf of the calcareous red algae *Corallina officinalis*. Among this turf, a variety of subdominant brown macroalgae (i.e. *Adenocystis utricularis*, *Hormosira banksii*, *Splachnidium rugosum*, *Scytothamnus australis*, *Ralfsia verrucosa*, *Colpomenia sinuosa*, *Gelidium caulacanthum*, *Stictosiphonia arbuscula* and foliose, green alga *Ulva* sp. were rated abundant to rare in cover (Table 3).

Although less abundant than both middle and high tidal elevations, a variety of sessile animals take advantage of the shelter and refuge provided from waves, heat, and predation by the algal canopy. In particular, limpets (*C. ornata*, *C. radians*, *Siphonaria zelandica*) and the snakeskin chiton *Sypharochiton pelleriserpentis* with a strong ability to cling to the rocks, were rated common to occasional, many returning to a home spot where their shell has adapted to fit the rock and provide a snug fit that offers protection from the elements (see Figure 6, bottom right).



Figure 5. *Hormosira banksii* and *Corallina officinalis* on the low shore.

### 3. Results and Discussion (Continued)



Figure 6. Topshells, limpets and chitons from the upper, mid and low eulittoral zones.

These species graze on the calcareous red algae *Corallina officinalis*. Two topshells (*Turbo smaragdus*, *Diloma aethiops*) were commonly observed, although many mobile topshells seek refuge from the sun in crevices during the day and are therefore not common on the open shore when the tide is out. Also among the algal turf, other invertebrate species present included *Chamaeosipho columna* - rare.



#### Fixed Quadrats

Results from the fixed quadrats at Flat Point are summarised in Tables 4 and 5, and Figures 6 and 7, with photos of each quadrat presented in Appendix 1. The principle purpose of repeat sampling fixed quadrats over time is to collect information on the stability of the mobile invertebrate and attached invertebrate and algal communities at representative shore heights. Because of the dynamic and often harsh rocky shore coastal environment, establishing a baseline of natural variability is vital if future changes are to be detected and interpreted. The baseline is designed to detect any long term vertical shift in the zonation pattern caused by sea level rise or changes in water quality (e.g. sea temperature or clarity) associated with climate change, and to evaluate impacts from introduced species, over-collection of shellfish, and from infrequent risks such as oil spills.

### 3. Results and Discussion (Continued)



Example of the rapid mobilisation of topshells from low tide refuges after being covered by water on the incoming tide.


Table 4 summarises richness, abundance and diversity measures for the three shore heights. A total of 19 species were recorded from quadrats, with fewer found at high and mid shore (10 and 7, respectively), compared to the lower shore (12). The diversity of both algae and mobile invertebrates was highest at the low shore, and the lowest on the upper shore. Note that this result only reflects species within the quadrats, and not the shore overall, as quadrat sampling excludes habitats such as crevices and rock pools which will support many additional species, while many mobile species move into shelter while the tide is out (see sidebar photo).

**Table 4. Summary of richness, abundance, and diversity indices for mobile invertebrates, sessile invertebrates, and macroalgae present in quadrats, Flat Point, 2016.**

Category	High Shore			Mid Shore			Low Shore		
	2016	2017	2018	2016	2017	2018	2016	2017	2018
Total number of species	5			6			10		
<b>MOBILE INVERTEBRATES (topshells, limpets, chitons)</b>									
RICHNESS (Number of species)	2			5			4		
ABUNDANCE (Mean number of individuals)	28.0			12.8			0.3		
DIVERSITY (Shannon Index)	0.04			0.77			0.35		
<b>SESSILE INVERTEBRATES (barnacles, mussels)</b>									
RICHNESS (Number of species)	3			1			0		
ABUNDANCE (Mean percentage cover)	94.3			31.7			-		
DIVERSITY (Shannon Index)	0.08			-			-		
<b>MACROALGAE</b>									
RICHNESS (Number of species)	0			0			8		
ABUNDANCE (Mean percentage cover)	-			-			66.2		
DIVERSITY (Shannon Index)	-			-			0.45		

Note: Low shore macroalgal percent cover values can exceed 100% because of overlapping algal growth.

Figure 7 shows that the biotic community present predictably groups quadrats relatively strongly, based on shore height. Such groupings also confirm that the individual sampling locations selected at each shore height are representative of each other. The similarity between the high and middle shore communities among quadrats (i.e. lack of distance between high and mid shore zones) reflects the relatively narrow tidal range with these sites being very similar in appearance and located close together on the shore.



**Figure 7. n-MDS plot showing the relationship among samples in terms of similarity in macroalgal and macroinvertebrate community composition for Flat Point rocky shore quadrats, Jan. 2016.**

The n-MDS plot (left) shows the 6 replicate samples at each of three shore heights, based on Bray Curtis dissimilarity and square root transformed data. The approach involves multivariate data analysis methods, in this case non-metric multidimensional scaling (n-MDS) using PRIMER-e version 6.1.16. The analysis basically plots the site and abundance data for each species as points on a distance-based matrix (a scatterplot ordination diagram). Points clustered together are considered similar, with the distance between points and clusters reflecting the extent of the differences. The interpretation of the ordination diagram depends on how well it represents actual dissimilarities (i.e. how low the calculated stress value is). Stress values greater than 0.3 indicate that the configuration is no better than arbitrary, and we should not try and interpret configurations unless stress values are less than 0.2.

### 3. Results and Discussion (Continued)

**Table 5. Raw data, mean number or percentage cover ( $\pm$ SE) and SACFOR rating of mobile invertebrates, sessile invertebrates, and macroalgae present in high, mid, and low shore quadrats, Flat Point, 2016.**

#### High Shore Quadrat Data

H2016	Scientific name	Common Name	Unit	Class	Quadrat						Total		
					1	2	3	4	5	6	Mean	SE	SACFOR
Limpets	<i>Cellana ornata</i>	Ornate limpet	#	ii	10	21	16	27	56	37	27.8	6.79	A
Limpets	<i>Cellana radians</i>	Tortoiseshell limpet	#	ii		1					0.2	-	R
Barnacles	<i>Chamaesipho columna</i>	Column barnacle	%	i	50	95	95	95	90	80	84.2	7.24	S
	<i>Epopella placata</i>	Ridged barnacle	%	i					1		0.2	-	R
Bivalves	<i>Xenostrobus neozelanicus</i>	Black mussel	%	i	5	15	15	10	5	10	10.0	1.83	O

#### Mid Shore Quadrat Data

M2016	Scientific name	Common Name	Unit	Class	Quadrat						Total		
					1	2	3	4	5	6	Mean	SE	SACFOR
Topshells	<i>Haustrum scobina</i>	Oyster borer	#	ii			1				0.2	-	R
	<i>Risellopsis varia</i>	Ridged periwinkle	#	ii				1			0.2	-	R
Limpets	<i>Cellana ornata</i>	Ornate limpet	#	ii	4	14	5	8	10	14	9.2	1.76	C
	<i>Cellana radians</i>	Tortoiseshell limpet	#	ii			1	6		1	1.3	1.18	F
Chitons	<i>Sypharochiton pelliserpentis</i>	Snakeskin chiton	#	ii			11	1			2.0	2.89	F
Barnacles	<i>Chamaesipho columna</i>	Column barnacle	%	i	40	35	40	35	25	15	31.7	4.01	C

#### Low Shore Quadrat Data

L2016	Scientific name	Common Name	Unit	Class	Quadrat						Total		
					1	2	3	4	5	6	Mean	SE	SACFOR
Topshells	<i>Diloma aethiops</i>	Grooved topshell	#	ii	1						0.2	-	R
Brown Algae	<i>Adenocystis utricularis</i>	Sea bladder	%	ii				5	5		1.7	-	O
	<i>Ralfsia verrucosa</i>	Tar spot/blood crust	%	i				1			0.2	-	R
	<i>Hormosira banksii</i>	Neptunes necklace	%	ii		2		3	2	8	2.5	1.17	O
	<i>Colpomenia sinuosa</i>	Oyster thief	%	ii		2	1	3	2		1.3	0.33	O
Green Algae	<i>Ulva lactuca</i>	Sea lettuce	%	ii	1				4	2	1.2	0.62	O
Red Algae	<i>Corallina officinalis</i>	Pink turf	%	ii	60	25	50	60	15	65	45.8	8.51	S
	<i>Gelidium caulacanthum</i>		%	ii		30		15	10	25	13.3	3.73	C
	<i>Stictosiphonia arbuscula</i>	Moss weed	%	ii	1						0.2	-	R



### 3. Results and Discussion (Continued)




Figure 8. Mean SACFOR rating for species present in 6 fixed quadrats in high, mid and low eulittoral zones.

### 3. Results and Discussion (Continued)



The monitoring of representative rocky shore habitats in Greater Wellington is vital if these highly valued and ecologically important ecosystems are to be managed effectively. Key physical variables such as sea temperature and wave forces can underpin a wide range of physiological and ecological processes, including altered species interactions, predation intensity, and dispersal and tolerances to thermal stress (Schiel 2011). These can be driven by natural changes in large scale events such as the El Niño/La Niña-Southern Oscillation, or by human impacts on global climate systems. In addition, coastal ecosystems are directly and often significantly affected by human use and development (over-collection of living resources and introduction of invasive species), as well as changes in land-use practices that in particular alter sediment and nutrient loadings.

Kelp communities comprise the dominant biogenic habitat along temperate rocky shores and loss of the three-dimensional algal community will likely result in a cascade of effects trending towards lower value, two-dimensional habitat, dominated by low-lying crusts and turfs, with subsequent adverse impacts on fish, invertebrate and algal sub-canopy communities. Because declines in algal habitat have been linked to degradation of water quality, increased sedimentation, increased nutrients, and contaminant discharges (e.g. Foster and Schiel 2010, Fong 2008), ensuring these stressors remain at a level the coastal environment can assimilate is clearly very important.

The initial monitoring results from Flat Point indicate a healthy and unpolluted rocky shore community. The risk from pathogens, sedimentation, eutrophication, and toxins is currently considered low, while a low-moderate risk is present, based on predicted accelerated sea level rise and temperature change.

Once the baseline monitoring is complete, data analysis will include a combination of PRIMER-e based analyses (Clarke & Gorley 2006), analysis of similarity (ANOSIM) to test for statistical differences in biotic assemblages among quadrats over time, and similarity percentage analysis (SIMPER) to identify taxa contributing most to multivariate differences among quadrats over time. SIMPER analysis will also enable identification of key indicator species, which will aid the intended development of condition ratings to characterise the status of the shore. Such ratings have not previously been attempted because current scientific knowledge of many NZ rocky shore species is scarce or incomplete. However, by focusing on measuring shifts in community composition, the presence or absence of key indicator species (including introduced plants and animals), as well as indicators of nutrient enrichment and sedimentation, it will be possible to develop appropriate condition ratings once the baseline monitoring is completed.

In addition, the scheduled 3 years of baseline monitoring will provide a robust measure of natural variation against which any future shift in vertical zonation on the shoreline or community composition can be assessed. It will also provide an invaluable benchmark for assessing the possible impacts from infrequent events such as oil spills or toxic algal blooms should they occur.

### 4. SUMMARY

The inaugural year of baseline rocky shore monitoring at Flat Point showed a healthy and unpolluted coastline supporting a collection of common rocky shore organisms present in a predictable shoreline zonation.

The zonation extended from a relatively low diversity, high shore intertidal community, dominated by barnacles, limpets and mussels through the mid shore barnacle dominated zone where topshells, limpets and chitons were also common, to the highest diversity low shore algal zone dominated by the extensively turfing, crustose red algae *Corallina officinalis*, and beneath low water by several fleshy brown macroalgae (e.g. *Carpophyllum flexuosum*, *Carpophyllum plumosum* and *Cystophora torulosa*).

## 5. MONITORING

Flat Point has been identified by Greater Wellington Regional Council as a priority for monitoring the effects of predicted accelerated sea level rise and temperature change, the over-collection of living resources, the introduction of invasive species, and impacts from excessive sedimentation, eutrophication, pathogens and toxins. It is recommended that monitoring continue as outlined below:

### **Rocky Shore Monitoring:**

- Continue the scheduled baseline monitoring at Flat Point in January 2017. After the 3 year baseline is completed, reduce monitoring to 5 yearly intervals or as deemed necessary based on rocky shore condition ratings (to be developed).


## 6. ACKNOWLEDGEMENTS

Thanks to Megan Oliver (GWRC) for her support and feedback on the draft report.


## 7. REFERENCES

- Airoidi, L. 2003. *The effects of sedimentation on rocky coast assemblages*. *Oceanogr. Mar. Biol.* 41, 161–236.
- Airoidi, L., Bacchiocchi, F., Cagliola, C., Bulleri, F., Abbiati, M. 2005. *Impact of recreational harvesting on assemblages in artificial rocky habitats*. *Marine Ecology Progress Series* 299: 55–66.
- Clarke, K.R., Gorley, R.N. 2006. *PRIMER-e v6: User Manual/Tutorial*. PRIMER-E, Plymouth.
- Foster, M.S., Schiel, D.R. 2010. *Loss of predators and the collapse of southern California kelp forests (?): alternatives, explanations and generalizations*. *Journal of Experimental Marine Biology and Ecology* 393, 59–70.
- Fong, P. 2008. *Macroalgal-Dominated Ecosystems, Chapter 20*. In: Capone, D.G., Bronk, D.A., Mullholland, M.R., Carpenter, E.J. (Eds) *Nitrogen in the Marine Environment 2nd Ed.*, Elsevier Inc, ISBN 978-0-12-372522-6. pp917-947.
- Hiscock, K. (Ed.) 1996. *Marine Nature Conservation Review: Rationale and methods*. Unpublished report by the Joint Nature Conservation Committee, Peterborough, U.K.
- Hiscock, K. 1998. *Biological monitoring of marine Special Areas of Conservation: a handbook of methods for detecting change*. Joint Nature Conservation Committee, Peterborough.
- MNCR. 1990. *Use of the Marine Nature Conservation Review SACFOR abundance scales*. JNCC [www.jncc.gov.uk/page-2684](http://www.jncc.gov.uk/page-2684).
- Robertson, B.M., Stevens, L. 2007. *Kapiti, Southwest, South Coasts and Wellington Harbour - Risk Assessment and Monitoring recommendations*. Prepared for Greater Wellington Regional Council. 60p.
- Schiel, D.R. 2011. *Biogeographic patterns and long-term changes on New Zealand coastal reefs: Non-trophic cascades from diffuse and local impacts*. *Journal of Experimental Marine Biology and Ecology* 400 (2011) 33–51.
- Taylor, D.I., Schiel, D.R. 2010. *Algal populations controlled by fish herbivory across a wave exposure gradient on southern temperate shores*. *Ecology* 91, 201–211.
- Tomanek, L., Helmuth, B. 2002. *Physiological ecology of rocky intertidal organisms: a synergy of concepts*. *Int. Comp. Biol.* 2002, 42:771-775.


# APPENDIX 1. DETAILED RESULTS

High Eulittoral	2016	2017	2018
QUADRAT 1  NZTM 1847960 East NZTM 5429657 North			
QUADRAT 2  NZTM 1847960 East NZTM 5429657 North			
QUADRAT 3  NZTM 1847958 East NZTM 5429655 North			
QUADRAT 4  NZTM 1847951 East NZTM 5429653 North			
QUADRAT 5  NZTM 1847950 East NZTM 5429651 North			
QUADRAT 6  NZTM 1847949 East NZTM 5429648 North			

# APPENDIX 1. DETAILED RESULTS (CONT.)

Mid Eulittoral	2016	2017	2018
<p>QUADRAT 1</p> <p>NZTM 1847960 East NZTM 5429657 North</p>			
<p>QUADRAT 2</p> <p>NZTM 1847558 East NZTM 5429655 North</p>			
<p>QUADRAT 3</p> <p>NZTM 1847957 East NZTM 5429654 North</p>			
<p>QUADRAT 4</p> <p>NZTM 1847951 East NZTM 5429655 North</p>			
<p>QUADRAT 5</p> <p>NZTM 1847950 East NZTM 5429651 North</p>			
<p>QUADRAT 6</p> <p>NZTM 1847948 East NZTM 5429649 North</p>			

# Appendix 1. DETAILED RESULTS (CONT.)

Low Eulittoral	2016	2017	2018
<p>QUADRAT 1</p> <p>NZTM 1847948 East NZTM 5429663 North</p>			
<p>QUADRAT 2</p> <p>NZTM 1847949 East NZTM 5429661 North</p>			
<p>QUADRAT 3</p> <p>NZTM 1847946 East NZTM 5429662 North</p>			
<p>QUADRAT 4</p> <p>NZTM 1847947 East NZTM 5429659 North</p>			
<p>QUADRAT 5</p> <p>NZTM 1847948 East NZTM 5429657 North</p>			
<p>QUADRAT 6</p> <p>NZTM 1847946 East NZTM 5429657 North</p>			